

Warmer temperatures reduce the vectorial capacity of malaria mosquitoes

Krijn P. Paaijmans, Simon Blanford, Brian H. K. Chan and Matthew B. Thomas

Biol. Lett. published online 21 December 2011
doi: 10.1098/rsbl.2011.1075

References

This article cites 19 articles, 3 of which can be accessed free

<http://rsbl.royalsocietypublishing.org/content/early/2011/12/15/rsbl.2011.1075.full.html#ref-list-1>

P<P

Published online 21 December 2011 in advance of the print journal.

EXiS Open Choice

This article is free to access

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right-hand corner of the article or click [here](#)

Advance online articles have been peer reviewed and accepted for publication but have not yet appeared in the paper journal (edited, typeset versions may be posted when available prior to final publication). Advance online articles are citable and establish publication priority; they are indexed by PubMed from initial publication. Citations to Advance online articles must include the digital object identifier (DOIs) and date of initial publication.

To subscribe to *Biol. Lett.* go to: <http://rsbl.royalsocietypublishing.org/subscriptions>

Warmer temperatures reduce the vectorial capacity of malaria mosquitoes

Krijn P. Paaijmans^{1,2,*}, Simon Blanford^{1,2,3},
Brian H. K. Chan^{1,2,3} and Matthew B. Thomas^{1,2}

¹Center for Infectious Disease Dynamics, ²Department of Entomology, and
³Department of Biology, Pennsylvania State University, University Park,
PA 16802, USA

*Author for correspondence (krijn@paaijmans.nl).

The development rate of parasites and pathogens within vectors typically increases with temperature. Accordingly, transmission intensity is generally assumed to be higher under warmer conditions. However, development is only one component of parasite/pathogen life history and there has been little research exploring the temperature sensitivity of other traits that contribute to transmission intensity. Here, using a rodent malaria, we show that vector competence (the maximum proportion of infectious mosquitoes, which implicitly includes parasite survival across the incubation period) tails off at higher temperatures, even though parasite development rate increases. We also show that the standard measure of the parasite incubation period (i.e. time until the first mosquitoes within a cohort become infectious following an infected blood-meal) is incomplete because parasite development follows a cumulative distribution, which itself varies with temperature. Including these effects in a simple model dramatically alters estimates of transmission intensity and reduces the optimum temperature for transmission. These results highlight the need to understand the interactive effects of environmental temperature on multiple host-disease life-history traits and challenge the assumptions of many current disease models that ignore this complexity.

Keywords: parasite infection; pathogen transmission; malaria risk; climate change; extrinsic incubation period; mortality

1. INTRODUCTION

The ecology of many vector–parasite/pathogen interactions is strongly influenced by environmental temperature [1]. Accordingly, it has been suggested that the dynamics and distribution of a range of vector-borne diseases, including malaria, dengue, viral encephalitis, schistosomiasis, Lyme disease and West Nile virus, could be impacted by climate change [1,2].

Predicting the extent of possible changes in disease patterns requires detailed understanding of how a suite of vector and parasite traits respond to temperature. However, in many cases the nature of the temperature-dependent relationships remains poorly defined. For example, vector competence, which describes the ability

of a vector to acquire, maintain and transmit a parasite/pathogen, is widely assumed to be temperature-insensitive. Yet, evidence from a limited number of studies indicates that vector competence can change with temperature [3–6].

Similarly, the standard degree-day models used to characterize the development of parasites/pathogens within the vector (defined as the extrinsic incubation period, or EIP) typically give a single value per temperature. During the EIP, pathogens go through very many replication cycles before migrating to the salivary glands where they can be transmitted to humans. The number of infectious mosquitoes (i.e. with pathogens in the salivary glands) in a mosquito cohort is expected to increase, from zero to the maximum number observed, over several days [7,8]. It is unclear how this distribution is affected by temperature, or how selecting the starting point, the median or endpoint of the distribution alters estimates of transmission intensity.

Here, using the rodent malaria *Plasmodium yoelii* and the Asian malaria vector *Anopheles stephensi*, we examine these standard assumptions and explore the implications for our understanding of the effects of temperature on disease transmission.

2. MATERIAL AND METHODS

Twenty-five mice (female C57Bl/6 laboratory mice, Charles River Laboratories) were inoculated with 10^5 *P. yoelii* parasites (clone 17XNL, from the WHO Registry of Standard Malaria Parasites, University of Edinburgh, UK). Four days after inoculation, approximately 2000 female *A. stephensi* mosquitoes (2–4 days old) were pooled in a single large cage and allowed to feed for 30 min on the anaesthetized mice. Post blood-feed, the females that took a full blood-meal were randomly distributed among four incubators (two cages per incubator) maintained at 20°C, 22°C, 24°C (the standard temperature for *P. yoelii* transmission) and 26°C \pm 1°C, with 90 \pm 5% relative humidity and 12 L : 12 D cycle photoperiod. Mosquitoes were fed *ad libitum* on 10 per cent glucose solution supplemented with 0.05 per cent paraaminobenzoic acid. Two and three days post blood-feed, mosquitoes were provided with egg laying bowls.

Mosquito salivary glands were dissected under a standard dissecting microscope, with 25 mosquitoes (randomly selected from the two cages) per temperature treatment per time-interval. We recorded if a mosquito harboured sporozoites in the salivary glands (hereafter referred to as ‘infectious mosquito’). During the first dissection time-point, midguts were also dissected to establish baseline malaria infection prevalence, i.e. the proportion of females with oocysts on their midgut. The number of oocysts per midgut was also recorded for each temperature treatment (square root transformed to meet normality assumptions). Daily inspection of the oocyst development for a small number of mosquitoes gave an indication of when oocysts were about to complete their development. Salivary glands were dissected on that day, and on at least 5 subsequent days to capture the cumulative sporozoite release. Additional mosquitoes were dissected over the subsequent one to two weeks to verify that maximum prevalence of infectious mosquitoes had been reached (hereafter, referred to as ‘maximum transmission prevalence’).

The cumulative change in proportion of infectious mosquitoes (b) over time was described by a logistic model $b = b_{\max}/(1 + e^{-k(t-t_m)})$, where b_{\max} is the upper asymptote (or the maximum transmission prevalence), t_m a constant and k the time at which the absolute increase in b is maximal. Best-fit logistic functions were used to estimate the maximum transmission prevalence, together with the EIP₁₀, EIP₅₀ and EIP₉₀ (i.e. the time to 10, 50 and 90% of the maximum prevalence) at a given temperature. To test if maximum infection prevalence varied significantly across temperature, we considered data points greater than EIP₉₀.

Transmission intensity was characterized using the standard formulation for vectorial capacity (C) for a single host–pathogen–vector system [9]:

$$C = \frac{ma^2 bcp^{\text{EIP}}}{-\ln(p)},$$

where m is the vector : human ratio; a vector biting frequency; p daily vector survival rate; bc , vector competence (a combination of b , the




Figure 1. Parasite development rates and vector competence over time at four temperatures (black squares, black dashed line, 20°C; black circles, black solid line, 22°C; grey squares, grey dashed line, 24°C; grey circles, grey solid line, 26°C). The data points indicate salivary infection prevalence at particular dissection times for the different temperature treatments. The lines represent best-fit logistic growth curves for each temperature as described in table 1. Inset shows the daily proportion of new infectious individuals in a mosquito cohort at the four different temperatures.

probability of a person becoming infected via a bite from an infectious vector, and c , the probability of a vector becoming infected by feeding on an infectious person); and EIP. Effects of temperature were explored using the empirical measures of b (maximum transmission prevalence), c (baseline infection prevalence) and EIP. There is very little information on the effects of temperature on the number of mosquito vectors per human so we set m arbitrarily to 1. Temperature has minimal effect on adult mosquito survival across the range of experimental temperatures explored [10]. However, since survival can vary substantially between malaria vector species [11] we chose two representative daily survival probabilities (0.8 and 0.9). Biting frequency, a , was described using the temperature-dependent function of Lardeux *et al.* [12].

3. RESULTS

Temperature did not affect baseline infection prevalence (mean of 91%; $\chi^2 = 3.57$, d.f. = 3, $p = 0.312$, hence c equals 0.91), nor the mean number of oocysts per midgut (221 ± 16 ; $F_{3,90} = 1.96$, $p = 0.125$). Increasing temperature did, however, increase parasite development rate ($1/\text{EIP}$), with the first infectious mosquitoes observed at days 15, 11, 9 and 8 at 20°C, 22°C, 24°C and 26°C, respectively (figure 1). Warmer temperatures also reduced the standard deviation in EIP among mosquitoes (inset figure 1 and table 1).

In contrast to development rates, increasing temperature did not simply increase vector competence. Maximum transmission prevalence was observed at 22°C (48%; figure 1 and table 1), and this prevalence was significantly higher than that at 24°C ($\chi^2 = 9.44$, d.f. = 1, $p = 0.002$). Prevalence at 24°C, in turn, was higher than that at 26°C ($\chi^2 = 24.24$, d.f. = 1, $p < 0.001$). The proportion of infectious mosquitoes was clearly reduced at 20°C but did not plateau by day 30 so was not analysed further.

The conventional approach of assuming temperature-independent vector competence (whereby $b = 0.24$, the

mean empirical value across temperature, and $c = 0.91$) predicted vectorial capacity to be highest at 26°C (figure 2). At this temperature, whether EIP was defined at the 10, 50 or 90th percentile of the distribution had little effect. At cooler temperatures, however, selecting the EIP₅₀ or EIP₉₀ reduced estimates of C relative to the EIP₁₀; the time taken to reach the much longer EIP₅₀ or EIP₉₀ at cooler temperatures means there are fewer mosquitoes alive to transmit.

Allowing temperature-dependent variation in vector competence changed these patterns considerably (figure 2). Maximum C is now predicted at 24°C, regardless of the daily mosquito survival probability, and exhibits a strong decline at lower and higher temperatures (i.e. compared with conventional estimates vectorial capacity is reduced 5.0-, 1-, 1.6- and 5.3-fold, at 20°C, 22°C, 24°C and 26°C, respectively). Moreover, in contrast to the standard predictions, use of EIP₅₀ or EIP₉₀ increased the estimates of C compared with those based on EIP₁₀; although it takes longer for 50 or 90 per cent of the infected mosquito population to become infectious, the much higher probability of being infectious at these later time points more than compensates for the increased cumulative daily mortality.

4. DISCUSSION

This study demonstrates marked effects of temperature on malaria parasite development rate and vector competence. As expected, warmer temperatures reduce the EIP. In addition, for a given ambient temperature the standard deviation of the EIP decreases as temperature (development rate) increases. We find that whether EIP is defined at an early (EIP₁₀) or late (EIP₉₀) stage of this distribution can have substantial impact on estimates of vectorial capacity. The observed effects of temperature on vector competence add further complexity. The strongly nonlinear relationship reduces the optimum temperature for transmission (vector competence trades-off against parasite development rate), and also reverses the relative influence of the EIP distribution for estimating vectorial capacity.

Our experiments used a rodent malaria and one species of mosquito and there is clearly a need to extend investigations to human malaria species and to other important vectors [13]. Nonetheless, the *A. stephensi*-*P. yoelii* system is considered a biologically realistic model [14] and there is no reason to believe that the temperature sensitivity of vector competence and the cumulative distribution in parasite development times are unique to this system.

The mechanisms behind the reduced vector competence at higher temperatures require further investigation but could include direct parasite mortality [15], reduced salivary gland invasion efficiency and sporozoite chemotaxis [16], and/or increased mosquito immune-related responses [17]. In addition, the transmission potential of individual mosquitoes across the EIP range needs to be determined as it is unclear whether mosquitoes at the EIP₁₀ and the EIP₉₀, for example, are equally infectious to a vertebrate host. Furthermore, our simplifying assumption setting the vector : host ratio (m) to 1 overlooks potentially complex effects of temperature on diverse mosquito traits such as




Figure 2. Mosquito vectorial capacity across temperature. Vectorial capacity estimated using either temperature-independent (black data points) or temperature-dependent (grey data points) measures of vector competence, and the EIP₁₀ (squares), EIP₅₀ (circles) or EIP₉₀ (triangles; i.e. the time to 10, 50 and 90% of the maximum prevalence at a given temperature) for a daily mosquito survival probability of (a) 0.8 and (b) 0.9.

Table 1. Parameters of the logistic growth model (and 95% CI) used to characterize the cumulative distribution of the number of mosquitoes that become infectious (*b*) over time at different temperatures. Parasite development times (EIP, in days) and proportion of infectious mosquitoes across temperature, when 10, 50 and 90% of the final infectious population is infectious are also shown.

	20°C	22°C	24°C	26°C
b_{\max}	9.7 (−6.5–26.0)	47.8 (46.7–49.0)	30.9 (29.6–32.1)	9.1 (5.0–13.3)
t_m	20.2 (19.4–21.0)	11.9 (6.5–17.4)	9.4 (5.9–12.9)	8.0 (−1.7E7–1.7E7)
k	0.3 (−14.7–15.3)	1.5 (1.0–2.1)	1.7 (1.2–2.2)	24.1 (−3.8E10–3.8E10)
R^2	0.501	0.954	0.962	0.539
EIP ₁₀	12.8	10.5	8.2	7.9
EIP ₅₀	20.2	11.9	9.4	8.0
EIP ₉₀	27.6	13.4	10.7	8.1
b_{10}	0.010	0.048	0.031	0.009
b_{50}	0.049	0.239	0.154	0.046
b_{90}	0.088	0.430	0.278	0.082

immature development and survival, adult survival and fecundity. Adding these effects could further alter vectorial capacity and suggests a need for additional experimentation, ideally under field conditions.

Overall, our results challenge current understanding of the effects of temperature on malaria transmission dynamics. We expect the effects to be robust across human malarias and possibly other vector-borne diseases. If so, the findings have significant implications for the various strategic modelling frameworks informing current disease control and eradication efforts [18,19], as they suggest that control at higher temperatures might be more feasible than currently predicted. The results also add complexity to studies investigating the possible effects of climate warming [20], as increases in temperature need not simply lead to increases in transmission.

This study was carried out in accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the NIH. The protocol was approved by the Animal Care and Use Committee of the Pennsylvania State University (Permit no.: 27452).

We thank Andrew Read, Sarah Reece and a further anonymous referee for constructive comments. This project

was part-funded by NSF (no. EF-0914384), NIH-NIAID (no. U19AI089676-01) and a grant with the Pennsylvania Department of Health using Tobacco Settlement Funds. The Department specifically disclaims responsibility for any analyses, interpretations or conclusions.

- 1 Githeko, A. K., Lindsay, S. W., Confalonieri, U. E. & Patz, J. A. 2000 Climate change and vector-borne diseases: a regional analysis. *Bull. WHO* **78**, 1136–1147.
- 2 Gage, K. L., Burkot, T. R., Eisen, R. J. & Hayes, E. B. 2008 Climate and vectorborne diseases. *Am. J. Prev. Med.* **35**, 436–450. (doi:10.1016/j.amepre.2008.08.030)
- 3 Stratman-Thomas, W. K. 1940 The influence of temperature on *Plasmodium vivax*. *Am. J. Trop. Med.* **20**, 703–715.
- 4 Siddons, L. B. 1944 Observations on the influence of atmospheric temperature and humidity on the infectivity of *Anopheles culicifacies* Giles. *J. Mal. Inst. India* **5**, 375–388.
- 5 Reisen, W. K., Fang, Y. & Martinez, V. M. 2006 Effects of temperature on the transmission of West Nile virus by *Culex tarsalis* (Diptera: Culicidae). *J. Med. Entomol.* **43**, 309–317. (doi:10.1603/0022-2585(2006)43[0309:EO TOTT]2.0.CO;2)
- 6 Kilpatrick, A. M., Meola, M. A., Moudy, R. M. & Kramer, L. D. 2008 Temperature, viral genetics, and

- the transmission of West Nile virus by *Culex pipiens* mosquitoes. *Plos Pathogens* **4**, e1000092. (doi:10.1371/journal.ppat.1000092)
- 7 Vaughan, J. A., Noden, B. H. & Beier, J. C. 1992 Population dynamics of *Plasmodium falciparum* sporogony in laboratory-infected *Anopheles gambiae*. *J. Parasitol.* **78**, 716–724. (doi:10.2307/3283550)
 - 8 Lambrechts, L., Paaijmans, K. P., Fansiri, T., Carrington, L. B., Kramer, L. D., Thomas, M. B. & Scott, T. W. 2011 Impact of daily temperature fluctuations on dengue virus transmission by *Aedes aegypti*. *Proc. Natl Acad. Sci. USA* **108**, 7460–7465. (doi:10.1073/pnas.1101377108)
 - 9 MacDonald, G. 1957 *The epidemiology and control of malaria*. London, UK: Oxford University Press.
 - 10 Craig, M. H., Snow, R. W. & Le Sueur, D. 1999 A climate-based distribution model of malaria transmission in sub-Saharan Africa. *Parasitol. Today* **15**, 105–111. (doi:10.1016/S0169-4758(99)01396-4)
 - 11 Kiszewski, A., Mellinger, A., Spielman, A., Malaney, P., Sachs, S. E. & Sachs, J. 2004 A global index representing the stability of malaria transmission. *Am. J. Trop. Med. Hyg.* **70**, 486–498.
 - 12 Lardeux, F. J., Tejerina, R. H., Quispe, V. & Chavez, T. K. 2008 A physiological time analysis of the duration of the gonotrophic cycle of *Anopheles pseudopunctipennis* and its implications for malaria transmission in Bolivia. *Malar. J.* **7**, 141. (doi:10.1186/1475-2875-7-141)
 - 13 Cohuet, A., Osta, M. A., Morlais, I., Awono-Ambene, P. H., Michel, K., Simard, F., Christophides, G. K., Fontenille, D. & Kafatos, F. C. 2006 *Anopheles* and *Plasmodium*: from laboratory models to natural systems in the field. *EMBO Rep.* **7**, 1285–1289. (doi:10.1038/sj.embor.7400831)
 - 14 Jaramillo-Gutierrez, G., Rodrigues, J., Ndikuyeze, G., Povelones, M., Molina-Cruz, A. & Barillas-Mury, C. 2009 Mosquito immune responses and compatibility between *Plasmodium* parasites and anopheline mosquitoes. *BMC Microbiol.* **9**, 154. (doi:10.1186/1471-2180-9-154)
 - 15 Kutz, S. J., Jenkins, E. J., Veitch, A. M., Ducrocq, J., Polley, L., Elkin, B. & Lair, S. 2009 The Arctic as a model for anticipating, preventing, and mitigating climate change impacts on host-parasite interactions. *Vet. Parasitol.* **163**, 217–228. (doi:10.1016/j.vetpar.2009.06.008)
 - 16 Akaki, M. & Dvorak, J. A. 2005 A chemotactic response facilitates mosquito salivary gland infection by malaria sporozoites. *J. Exp. Biol.* **208**, 3211–3218. (doi:10.1242/jeb.01756)
 - 17 Hillyer, J. F., Barreau, C. & Vernick, K. D. 2007 Efficiency of salivary gland invasion by malaria sporozoites is controlled by rapid sporozoite destruction in the mosquito hemocoel. *Int. J. Parasitol.* **37**, 673–681. (doi:10.1016/j.ijpara.006.12.007)
 - 18 Tatem, A. J., Smith, D. L., Gething, P. W., Kabaria, C. W., Snow, R. W. & Hay, S. I. 2010 Ranking of elimination feasibility between malaria-endemic countries. *Lancet* **376**, 1579–1591. (doi:10.1016/S0140-6736(10)61301-3)
 - 19 The malERA Consultative Group on Modeling. 2011 A research agenda for malaria eradication: modeling. *PLoS Med.* **8**, e1000403. (doi:10.1371/journal.pmed.1000400)
 - 20 Parham, P. E. & Michael, E. 2010 Modeling the effects of weather and climate change on malaria transmission. *Environ. Health Perspect.* **118**, 620–626. (doi:10.1289/ehp.0901256)